



T1 measurements identify extracellular volume expansion in a genotyped hypertrophic cardiomyopathy population with and without left ventricular hypertrophy

Citation

Abbasi, Siddique, Ravi Shah, Tomas G Neilan, Bobby Heydari, Yucheng Chen, Michael Jerosch-Herold, Raymond Y Kwong, and Carolyn Ho. 2013. T1 measurements identify extracellular volume expansion in a genotyped hypertrophic cardiomyopathy population with and without left ventricular hypertrophy. *Journal of Cardiovascular Magnetic Resonance* 15(Suppl 1): 021.

Published Version

doi:10.1186/1532-429X-15-S1-021

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:11235263>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)



ORAL PRESENTATION

Open Access

T1 measurements identify extracellular volume expansion in a genotyped hypertrophic cardiomyopathy population with and without left ventricular hypertrophy

Siddique Abbasi^{1*}, Ravi Shah^{1,2}, Tomas G Neilan^{1,2}, Bobby Heydari¹, Yucheng Chen¹, Michael Jerosch-Herold¹, Raymond Y Kwong¹, Carolyn Ho¹

From 16th Annual SCMR Scientific Sessions
San Francisco, CA, USA. 31 January - 3 February 2013

Background

Myocardial fibrosis is a hallmark of hypertrophic cardiomyopathy (HCM) and may contribute to arrhythmias and heart failure. Sarcomere mutations appear to induce profibrotic changes before left ventricular hypertrophy (LVH) develops. Measuring T1 relaxation times with contrast cardiac magnetic resonance (CMR) allows robust quantification of the cardiac extracellular volume (ECV) and non-invasive assessment of diffuse myocardial fibrosis.

Methods

A genotyped HCM population underwent contrast CMR with measurement of T1. Subjects included sarcomere

mutation carriers with LVH (G+/LVH+, n = 37) and without LVH (G+/LVH-, n = 30); HCM patients without mutations (sarcomere-negative HCM, n = 11); and mutation-negative healthy controls (n = 10). Concurrent echocardiography and serum biomarkers of collagen synthesis, hemodynamic stress, and myocardial injury were available in a subset.

Results

Late gadolinium enhancement (LGE) was present in >60% of overt HCM patients but absent from G+/LVH- subjects. Compared to controls, ECV was increased in patients with overt HCM, as well as G+/LVH- mutation

Table 1 CMR Metrics of Fibrosis and Serum PICP levels

	Control N=10	p-value* Control vs Preclinical	Preclinical HCM N=30	p-value* Preclinical vs Overt	Overt HCM N=37	p-value* Control vs Overt
ECV (range)	0.26 ± 0.01 (0.24-0.31)	<0.001	0.33 ± 0.01 (0.23-0.38)	<0.001	0.37 ± 0.01 (0.31-0.51)	<0.001
ECV, excluding LGE	0.26 ± 0.01	<0.001	0.33 ± 0.01	<0.001	0.37 ± 0.01	<0.001
LGE present, % (#yes / #no)	0% (0/10)		0% (0/29)		78.4% (29/8)	
LGE, g	0	0.8	0	0.004	16.7 ± 4.9	0.003
LGE, % LV mass	0	0.7	0	0.0006	7.76 ± 1.96 (0.9-44.6%)	0.001
PICP, µg/L	63.08 ± 4.19	<0.001	83.81 ± 4.39	0.035	108.04 ± 9.81	<0.001

Data are presented as means adjusted for family relations and age ± standard error *p values <0.017 are statistically significant and adjusted for age and family relations ECV = extracellular volume; LGE = late gadolinium enhancement; PICP = carboxy-terminal propeptide of procollagen type I

¹Department of Cardiovascular Medicine, Brigham and Women's Hospital/Harvard Medical School, Boston, MA, USA

Full list of author information is available at the end of the article




Figure 1 ECV is significantly increased in HCM sarcomere mutation carriers both with and without LVH. Compared to normal controls, ECV was 25% in G+/LVH- subjects and 42% higher in G+/LVH+ HCM patients ($p<0.001$ for all comparisons to controls). ECV was further increased by 13% in overt HCM compared to G+/LVH- subjects ($p=0.0003$).

carriers (ECV, 0.37 ± 0.01 , 0.33 ± 0.01 , 0.26 ± 0.01 in G⁺/LVH+, G⁺/LVH-, controls, respectively, $P<0.001$). ECV was correlated with NT-proBNP levels ($r=0.61$, $P<0.001$) and global E' velocity ($r=-0.48$, $P<0.001$). ECV and LGE were higher in sarcomeric HCM than sarcomere-negative HCM.

Conclusions

Myocardial ECV is increased not only in sarcomere mutation carriers with overt HCM, but also those with normal LV wall thickness. These data provide further evidence that fibrotic remodeling is triggered early in disease pathogenesis. Assessing ECV may help characterize the development of HCM-induced myocardial fibrosis, ultimately supporting novel strategies to modify disease by targeting interstitial fibrosis.

Funding

This work was supported by grants from the National Institutes of Health (CYH and RYK).

Author details

¹Department of Cardiovascular Medicine, Brigham and Women's Hospital/Harvard Medical School, Boston, MA, USA. ²Cardiology, Massachusetts General Hospital, Boston, MA, USA.

Published: 30 January 2013

doi:10.1186/1532-429X-15-S1-O21

Cite this article as: Abbasi et al.: T1 measurements identify extracellular volume expansion in a genotyped hypertrophic cardiomyopathy population with and without left ventricular hypertrophy. *Journal of Cardiovascular Magnetic Resonance* 2013 **15**(Suppl 1):O21.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

