



P01-01. The Blood Transcriptional Response to Early Acute HIV Infection is Transient and Responsive to Antiretroviral Therapy

Citation

Skinner, J. A., M. Sharma, N. Baldwin, B. Lemoine, D. Blankenship, E. B. De Vol, A. Mejias, et al. 2009. P01-01. The blood transcriptional response to early acute HIV infection is transient and responsive to antiretroviral therapy. *Retrovirology* 6(Suppl 3): P1.

Published Version

doi:10.1186/1742-4690-6-S3-P1

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:4872637>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Poster presentation

Open Access

P01-01. The blood transcriptional response to early acute HIV infection is transient and responsive to antiretroviral therapy

JA Skinner*¹, M Sharma¹, N Baldwin¹, B Lemoine¹, D Blankenship², EB De Vol², A Mejias³, O Ramilo³, M Cohen⁴, N Letvin⁵, D Goldstein⁶, K Soderberg⁷, TN Denny⁶, K Shianna⁶, A McMichael⁸, B Haynes⁷, J Banchereau¹ and D Chaussabel¹

Address: ¹Baylor Institute for Immunology Research, Dallas, TX, USA, ²Baylor Health Care System, Dallas, TX, USA, ³Dept. of Pediatrics, UTSSW, Dallas, TX, USA, ⁴University of North Carolina, Chapel Hill, NC, USA, ⁵Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA, USA, ⁶Duke Institute for Genome Sciences and Policy, Duke University, Durham, NC, USA, ⁷Duke University School of Medicine, Durham, NC, USA and ⁸Weatherall Institute of Molecular Medicine, Oxford, UK

* Corresponding author

from AIDS Vaccine 2009
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, **6**(Suppl 3):P1 doi:10.1186/1742-4690-6-S3-P1

This abstract is available from: <http://www.retrovirology.com/content/6/S3/P1>

© 2009 Skinner et al; licensee BioMed Central Ltd.

Background

Systemic events in acute HIV infection (AHI) are associated with disease severity and progression to AIDS. Timely identification of AHI patients has posed a significant challenge to identifying the underlying mechanisms driving these events. The aim of this study is to elucidate these pathways by characterizing the genome-wide transcriptional signature expressed by whole blood during early AHI.

Methods

Longitudinal whole blood samples from ART treated and untreated patients from both the United States (n = 16) and Africa (n = 16) were collected at study enrollment and weeks 1, 2, 4, 12, and 24. AHI and non-infected controls were analyzed using Illumina HT-12 microarrays. Both gene and module level analysis were conducted to identify biologic pathways active in AHI.

Results

Nineteen annotated and 24 undefined transcriptional modules constitute a robust transcriptional signature that collectively distinguished early AHI patients from non-infected controls. The activity of transcriptional modules related to interferon, cell cycle, cytotoxic, and mitochond-

drial responses were significantly increased in AHI patients. At study enrollment, the intensity of this signature was not correlated with viral load and exhibited heterogeneity between patients. Association between viral load and signature intensity was found over time. However, three patients exhibited little change in transcriptional activity despite high viral loads. When compared to acute RSV and Influenza infections, only interferon signatures were conserved across all three infections while cell cycle, cytotoxic, and mitochondrial responses were unique to AHI. The AHI signature of untreated patients regressed to non-infected control levels by 12–24 weeks post enrollment. The initiation of ART accelerated the dissipation of this signature, returning the core AHI signature to normal levels within 4 weeks.

Conclusion

The whole blood AHI transcriptional signature is unique, transient, and capable of classifying individual responses to infection. This signature is responsive to ART and contains pathways with both defined and novel associations with HIV infection.